



Atmospheric Photochemistry Studies of Pollutant Emissions from Transportation Vehicles Operating on Alternative Fuels

Subcontractor

University of North Carolina at Chapel Hill

Principal Investigator

Harvey E. Jeffries

University of North Carolina at Chapel Hill
Chapel Hill, NC 27599
(919) 966-7312

DOE Project Manager

Michael Gurevich

U.S. Department of Energy
CE-332, MS 6A-116/Forrestal
1000 Independence Avenue, SW
Washington, DC 20585
(202) 586-6104

NREL Technical Monitor

Brent Bailey

NREL

1617 Cole Boulevard
Golden, CO 80401-3393
(303) 275-4468

Subcontract Number

XAP-3-11153-01

Performance Period

9/93-9/95

NREL Subcontract Administrator


Scott Montgomery (303) 275-3193

Objective

To improve our ability to predict the changes in urban ozone (O_3) that result from the widespread use of alternative fuels in automobiles. The entire range of fuels—alcohol fuels, natural gas fuels, and reformulated gasoline (RFG)—is of interest.

Approach

The smog chamber is a chemical environment designed to simulate the complex volatile organic compound



The University of North Carolina outdoor smog chamber

(VOC) compositions of urban atmospheres. We conducted outdoor smog chamber experiments using complex VOC mixtures that arise from the use of blended ethanol (EtOH), compressed natural gas (CNG), and liquefied petroleum gas (LPG) as vehicle fuels. Advanced analytical methods are used to identify and quantify previously unknown or undetected reaction products. These results will then be used to test and improve the formulations of our current photochemical reaction mechanisms, and with Urban Airshed Model (UAM) simulations, to help answer questions concerning the effect of urban O_3 from the use of alternative fuels.


Accomplishments

Twenty-five smog chamber experiments were conducted for this study. Results indicate a substantial benefit from the use of CNG and LPG in a synthetic urban air mixture (SynURB) background. EtOH use did not result in an ozone reduction. Under some conditions, it increased ozone production. We also reformulated a commonly used airshed reaction mechanism to fit the chamber observations more accurately, and used this improved mechanism in UAM simulations of air quality under alternative fuel use scenarios.



Future Direction

A final report is expected to be published by February 1997.



Smog chamber results comparing ozone (O_3) formation from natural gas and from Industry Average Gasoline (IAG)

Publications

Borbridge, L. 1995. "ADual Derivatization Technique for the Determination of Unsaturated and Epoxy-Carbonyls in Air Using Gas Chromatography/Mass Spectrometry." University of North Carolina, August.

Borbridge, L.; H. Jeffries; J. Yu. 1995. "A Dual Derivation Technique for the Determination of Unsaturated and Epoxy-Carbonyls in Air Using Gas Chromatography/Mass Spectrometry," submitted to Analytical Chemistry.

Jang, J.C.C.; H.E. Jeffries; D. Byun; J.E. Pleim. 1995. "Sensitivity of Ozone to Model Grid Resolution, Part II," *Atmospheric Environment*, in press.

Lo, C.Y.S. 1995. "Process Composition Analysis of Urban Airshed Model Predictions." University of North Carolina, August.

Rigsbee, C. 1995. "The Hydroxy Radical Initiated Oxidation of C2-C4 Olefins: Detection of Carbonyl Compounds." University of North Carolina, August.

Yu, J.; H. Jeffries; R.M. LeLacheur. 1995. "Identifying

Airborne Carbonyl Compounds in Isoprene Atmospheric Photooxidation Products by Their PFBHAoximes Using Gas Chromatography/Ion Trap Mass Spectrometry," *Environmental Science & Technology*, Vol. 29, No. 8.

Jeffries, H; K. Sexton; J. Yu. 1996. "Atmospheric Photochemical Studies of Pollutant Emissions from Transportation Vehicles Operating on Alternative Fuels." Draft report to the National Renewable Energy Laboratory. Subcontract XAT-3-11153-01, April.